



MEETING ABSTRACT

Open Access

Varying modulation of HTLV-1 LTR activity by BAF complexes

Rachel Van Duyne^{1,2}, Irene Guendel¹, Nazly Shafagati¹, Kylene Kehn-Hall¹, Rebecca Easley¹, Zachary Klase³, Sergei Nekhai⁴, Mudit Tyagi¹, Fatah Kashanchi^{1,2*}

From 15th International Conference on Human Retroviruses: HTLV and Related Viruses
Leuven and Gembloux, Belgium. 5-8 June 2011

Chromatin remodeling is a rapidly emerging field with critical implications for the control of viral gene expression, especially for viruses with integrated genomes, such as HTLV-1. Recent observations indicate that there are as many as eight different BRG1 containing chromatin remodeling complexes highlight the advancement in the field, but also the necessity for future study especially when looking at viral infections. In the current study we focused on few of the Baf subunits that are common to most SWI/SNF complexes. We find that at least three Bafs, Baf53, Baf155 and Baf170, are highly regulated in HTLV-1 infected cells. Along these lines others have shown that depletion of Baf53 leads to the expansion of chromosome territories and decompaction of the chromatin. Here we show that there are clear varying differences between the Baf subunits after viral infection. These subunits also co-elute in different places from a sizing column and one particular form, Baf53 may be phosphorylated in HTLV-1 infected cells. Normally Baf53 is a suppressive complex and knock down experiments show increased level of virus gene expression from transfected or chronically infected cells.

Published: 6 June 2011

doi:10.1186/1742-4690-8-S1-A180

Cite this article as: Van Duyne et al.: Varying modulation of HTLV-1 LTR activity by BAF complexes. *Retrovirology* 2011 **8**(Suppl 1):A180.

Author details

¹George Mason University, Department of Molecular and Microbiology, National Center for Biodefense and Infectious Diseases, Manassas, VA, 20110, USA. ²The George Washington University Medical Center, Department of Microbiology, Immunology, and Tropical Medicine, Washington, DC, 20037, USA. ³National Institutes of Health, Molecular Virology Section, Laboratory of Molecular Microbiology, NIAID, Bethesda, Maryland, 20892, USA. ⁴Howard University, Department of Microbiology, Center for Sickle Cell Disease, Department of Medicine, Washington DC, 20060, USA.

* Correspondence: fkashanc@gmu.edu

¹George Mason University, Department of Molecular and Microbiology, National Center for Biodefense and Infectious Diseases, Manassas, VA, 20110, USA

Full list of author information is available at the end of the article

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

